



COLOMBIA OFF-THE-BEATEN-TRACK TOUR REPORT 2024

17 January - 5 / 12 February 2024

Trevor Ellery

We started the tour with an early departure from Bogota, so as to avoid the traffic and journeyed north through the Cundinamarca and Boyacá altiplanos, along the spine of the Eastern Andes. This area has been heavily populated for many years and natural habitat was scarce, though we did manage a few of the more common open country species, including Great Thrush, Eared Dove, Rufous-collared Sparrow and Western Cattle Egret. As we wended our way further into the more remote and mountainous regions, we began to see some remnant oak forest patches clinging to the steep slopes. Once they became more extensive, we stopped and began searching on the off chance that our main target, the endemic Colombian Mountain Grackle, may still be present. While we failed to find the Grackle, we did locate a selection of commoner Andean species, including White-throated Tyrannulet, Cinnamon Flycatcher, Brown-capped Vireo, Ashy-throated Chlorospingus, Golden-fronted Whitestart (of the white spectacted, ornatus subspecies), Scarlet-bellied Mountain Tanager, Longuemare's Sunangel and some nice White-tipped Swifts. We arrived in the arid Chicamocha canyon in the late morning and immediately visited a new Hummingbird garden. We found that the once tricky endemic Chestnut-bellied Hummingbird, was abundant at these feeders. Other species included the endemic Indigo-capped Hummingbird, Rufous-tailed Hummingbird, Black-throated Mango and White-bellied Woodstar. The surrounding scrub and gardens held Yellow-backed and Baltimore Orioles, Tropical Parula, Bicoloured wren and both Scrub and Tooth-billed Tanagers. We then dropped down to the lower valley for lunch and a post lunch stroll added a vocal pair of the endemic Apical Flycatcher. With the heat of the day subsiding, it was time to climb back to slightly higher elevations and it only took a very brief search to locate our fourth endemic of the day, a very obliging Niceforo's Wren. It was then decided to climb up to the lower reaches of the Oak forest above the valley, once again in search of in search of the Mountain Grackle. We failed to locate any Grackles but we did find White-collared Swift, Buff-tailed Coronet, our first Inca Jays and a Pale-bellied Tapaculo.

The following morning, we left before dawn and climbed to the ridge line, where we quickly located a couple of calling White-throated Screech Owls, which showed well. We then spent much of the morning searching for Mountain Grackle without success. A change of tactics was needed, so we crossed over the ridge and descended to the thick Oak forest on the other side. This almost immediately produced results, as a garrulous party of noisy Mountain Grackles moved into view. The Grackles were to prove much commoner on this side of the mountain and we saw multiple groups during the morning, which we estimated totalled a remarkable forty-eight individuals. Other species we encountered including a magnificent pair of Powerful Woodpeckers and a couple of brief sightings of Rusty-faced Parrot, with eight birds passing overhead and a pair flying out from directly beneath us, when scanning from a lookout. A distant but still impressive male Red-hooded Tanager was an unexpected bonus. We continued to amass a steady list of Andean species with Speckled Hummingbird, Tyrian Metaltail, Smoky-brown and Crimson-mantled Woodpeckers, Montane Woodcreeper, Black-capped Tyrannulet, Smoke-coloured Pewee, Rufous-breasted Chat-Tyrant, Yellow-bellied Siskin, Masked Trogon, Beryl-spangled Tanager and Mountain Cacique. The number of overwintering Blackburnian Warblers was also notable, with it being the most abundant species in any mixed flock and seeming to be present at a higher density in these Oak forests, than anywhere else in the Andes. Clearly these highly threatened and rather unique forests, are of great importance for boreal migrants, as well as a small suite of Colombian endemics.

Our third day above the Chicamocha canyon was spent searching for Gorgeted Wood Quail but despite much effort, we could find no sign. We did add Chestnut-crowned Antpitta, Glossy-black Thrush and Slate-throated Whitestart, along with many of the species that we had seen the previous day. We then had a long journey back to Bogota, with a couple of brief roadside stops adding Brown-backed Chat-Tyrant and White-tailed Kite.

Our fourth day began with an early morning flight to the bustling city of Valledupar in northern Colombia. We arrived at the oppressive heat of this narrow valley, wedged between the Santa Marta and Perija mountain ranges. We quickly met our jeeps and were whisked up into the cooler foothills of the Perija Mountains. Our first target was the endemic Perija Brushfinch and after a short search, we enjoyed good views of a rather skulking individual. We also noted an obliging Moustached Puffbird and picked up our first Golden-crowned Warblers. While plenty of commoner species were present at the base of the mountains, we continued to ascend in search of specialities. At our next stop we staked out some flowers in the garden of a friendly campesino. While he was preparing us Coffee from his Finca, a lovely restricted range Rufous-shafted Woodstar appeared and danced around the orange blossoms in front of us. We then continued up the mountain and stopped to try for Maroon-chested Ground Dove, an erratic and nomadic species, which had recently been found in the Perija. Despite much effort we were only able to hear and not see this enigmatic species. Despite this disappointment we carried on birding and soon located our next endemic, with a Perija Tapaculo showing well. Other species that we found on the upper slopes included Andean Guan, a brief Black-and-chestnut Eagle, an obliging White-throated Toucanet and a Rufous Spinetail (of the Perija endemic *munoztebari* subspecies, which is a potential split). Further birding produced Spectacled Tyrannulet, Streak-throated Bush Tyrant, Oleaginous Hemispingus, Black-crested Warbler (a vocally distinct form) and Common Chlorospingus (of the Perija endemic *ponsi* subspecies). We arrived at our lodge in the late afternoon, where the feeders were busy with more Longuemare's Sunangels and both Lesser and Sparkling Violetears. The seed feeders meanwhile attracted the Perija endemic Black-fronted Brushfinch and the striking *fumidus* race of Slaty Brushfinch. We ended the day with a vocal male Band-winged Nightjar briefly flying around at dusk.

The following morning, we ascended high above the Lodge to the Paramo of Sabana Rubia, where we were to search for many of the key Perija endemics. Soon after dawn we located a vocal and eventually showy Perija Thistletail. We followed this with several endemic Perija Metaltails, with individuals of both sexes showing well. We did have to be careful though not to confuse them with the *districta* race Tyrian Metaltails, which were also present (and which is also a potential split). The recently split Perija Antpitta proved trickier but we all eventually obtained reasonable views of this skulking species. Other avian highlights included Lacrimose Mountain Tanagers (of the endemic and distinctive *pallidorsalis*, which is yet another potential split) and an obliging Andean Pygmy Owl. One of the toughest endemics in the Perija is the recently split Perija Starfrontlet but we managed to see three, dashing about on the Paramo edge and we even obtained some record shots of a young male. We also added Andean Condor, Black-chested Buzzard-Eagle White-browed Spinetail, Red-crested Cotinga, Rufous-breasted Chat-Tyrant, Mountain Wren and a couple of Chestnut-breasted Chlorophonias. Mixed flocks held Hooded Mountain and Blue-capped Tanagers, Pearled Treerunner and Blue-backed Conebills.

The following day we made a slow descent from the Perija with the upper slopes producing Rufous-crowned Tody-Flycatcher, Yellow-bellied Chat-Tyrant, Buff-breasted Mountain Tanager, Plushcap and Coppery Emerald. The lower slopes also held plenty of new species for the trip, including Streak-headed Woodcreeper, Pale-eyed Pygmy Tyrant, Greenish Elaenia, Whiskered, Rufous-breasted and Rufous-and-white Wrens, Golden-winged Sparrow and Chestnut-capped Warbler. A fine selection of boreal migrants featured with Black-and-

white, Tennessee and Mourning warblers, American Redstart, Yellow-throated Vireo and Rose-breasted Grosbeak. We also logged a confiding Blue-black Grosbeaks, while new Tanagers included Black-faced, Crimson-backed, Blue-capped, Fawn-breasted and Black-headed. A late evening visit to the dry forests of the Los Besotes reserve was fairly quiet, with the extreme drought that Colombia was experiencing, possibly affecting the avian activity at this site. We did locate a male Pauraque, both Red-billed Emerald and Shining-green Hummingbird, Brown-throated Parakeet, Caribbean Hornero, Northern White-fringed Antwren and our first Scrub Greenlets. A final surprise was a pair of Double-striped Thick-knees, illuminated in the headlights on the drive out.

The next morning, we returned to Los Besotes and after much searching, we located a couple of Venezuelan Flycatchers. The forest and scrub were more productive in these early cooler hours and we also managed to find Crested Bobwhites, Scaled and Common Ground Doves and Plain-brown Woodcreeper. A hat trick of Antshrikes featured Black-crested, Black-backed and Black-crowned, which were all seen well. We also added a few species whose distribution creeps down from the nearby Guajira peninsula, including White-whiskered Spinetail, Glaucous Tanager and both Pale-tipped and Slender-billed Inezias. It was very productive for Flycatchers with Ochre-ored and Yellow-olive Flatbills, Forest Elaenia, Great-crested and Brown-crested Flycatchers and a surprise Fuscous Flycatcher. A few further commoner species included our first Yellow Orioles and Rufous-browed Peppershrike. We then made the transfer south to La Jagua del Ibrico. The main road was blocked by a local protest, so we had to extract our luggage from one van, walk across the roadblock and then connect with another van once we had passed it, an interesting experience! Upon arrival at La Jagua we climbed into some jeeps and headed back up into the mountains. Our destination was a remote Finca, where we hoped to connect with another Perija endemic. We were extremely fortunate, as almost immediately upon arrival we found a small group of the truly stunning Perija Parakeet, feeding in a bush just behind the kitchen! With our main target in the bag and having enjoyed some superlative views, it was time for lunch. We followed this with some afternoon birding, where we added Sooty-capped Hermit, Long-billed Gnatwren, and Cocoa Woodcreeper.

The following morning, we left early for the drive to the Bushbird Reserve, arriving at this rather remote location in the mid-morning. Some initial searching for the Bushbird produced either heard only or very brief glimpses. We did though connect with a range of other key species including Grey-throated Warbler, Stripe-breasted Spinetail and a wonderful Orange-billed Nightingale-Thrush. Towards the late afternoon we birded along the road, where we found some smart Chestnut-bellied Thrushes, the local Moustached Brushfinch and towards dusk, a nice group of Band-tailed Guans.

The following morning, we decided to try a different area and were rewarded with superb views of a pair of Recurve-billed Bushbirds, with the male being especially obliging. We also added our final key target, with a skulking Klages's Antbird showing well. Commoner species included Golden-bellied and Dusky-capped Flycatchers, Speckled Tanager and Plain Antvireo. It was then time to make the long drive to the Cerulean Warbler reserve, with a lunch stop in the Magdalena Valley adding a few commoner species. It was the final part of the drive that was to prove most productive though. Our first brief stop produced a nice male Jet Antbird and then, as we began to climb into the Cacao plantations, further judicious pauses allowed us to add Double-banded Greytail, Barred Puffbird, Violet-bellied Hummingbird, Plain-coloured Tanager, White-eared Conebill and Yellow-tufted Dacnis. We arrived at the lodge just in time to be shown a surprise, bonus roosting Black-and-white Owl! We were based in the Cerulean Reserve for several nights and we were to spend much of the next two days, on the trails in the lovely subtropical forest above the reserve. A new access track meant that it was possible to now drive most of the way up to the forest, which gave us more flexibility and we interspersed long hikes in the forest, with some birding at the lodge and in the surrounding Coffee and Cacao plantations.

Our first early morning visit to the forest produced a surprise Rufous-bellied Nighthawk, perching on fence posts near the trail entrance and we also managed to not just hear but also see a Highland Tinamou, scurrying down the trail. The Hummingbird feeders in the forest hosted the endemic Black Inca along with Green-crowned and Fawn-breasted Brilliants and White-booted Racket, while the seed feeder allowed good views of Ruddy Quail-Dove and Chestnut-capped Brushfinch. We stayed out past dusk on one night and managed superlative views of a very confiding Cinnamon Screech Owl. We also spent time working the flocks and searching the understorey in the forest and this proved very productive. We found Collared Trogon, Uniform Antshrike, Slaty Antwren, the endemic Parker's Antbird and a skulking pair of Blue-lore Antbirds. Ranging through the canopy were Rufous-rumped Antwren, Brown-billed Scythebill, Olive-backed Woodcreeper, Montane and Buff-fronted Foliage-gleaners, Variegated Bristle Tyrant, Plumbeous-crowned Tyrannulet and Rufous-naped Greenlet. We also enjoyed a couple of Chestnut-crowned Gnatcatchers and after much effort the endemic Magdalena Tapaculo. Further avian delights included Subtropical Cacique, Golden-winged Warbler, White-winged Becard, Metallic-green and Golden Tanagers and Blue-winged Mountain Tanager. A White-bellied Antpitta was also glimpsed after a very patient wait. When not in the forest we were invariably to be found in the vicinity of the lodge. The busy hummingbird feeder held Brown Violetear, White-necked Jacobin and Short-

tailed Emerald along with a plethora of commoner species. Other species in the grounds included both Spot-breasted and Lineated Woodpeckers and endemic Colombian Chachalacas, while a Black Hawk-Eagle was noted drifting overhead! We also spent an early morning staking out a favoured tree and duly enjoyed good scope views of the endemic Turquoise Dacnis, while a few stop on the drive up to main reserve, allowed good views of Black-headed Brushfinch and an outrageous six Cerulean Warblers!

Our final morning was spent slowly birding our way out of the San Vicente area but not before we had added a confiding Mottled Owl in the lodge grounds. We descend once again to the mid elevations on the mountain and a busy morning produced Collared Aracari, a very obliging Yellow-browed Shrike-Vireo, Laughing Falcon, Spectacled Parrotlet, Black-headed Tody-Flycatcher, Golden-rumped Euphonia and many commoner species. We then made the journey down the Magdalena valley and into the Paujil reserve, with some birding stops enroute. The long drive into Paujil was enlivened by our first Northern Screamers and the wetlands also held Green Ibis, Rufescent Tiger-Heron and White-headed Marsh Tyrant. The verdant vegetation along the track side produced Plain-breasted Ground Dove, Grey Seedeater and Thick-billed Seedfinch. We arrived at the lodge at dusk and a highlight of the evening was a Choco Screech Owl, which ended up in the forest guard's house. This necessitated an early wake up call for many of the group, who had only just climbed into bed! The Owl gave fantastic views, before it was captured and released back out into the night.

The following day we had a full day at the Paujil Reserve, where the main highlight was of course the very obliging Curassows. We had a spectacular twelve birds wandering around the Lodge, where they are now very habituated. We also enjoyed Blue-chested Hummingbird, Stripe-throated Hermit, a couple of Grey-cowled Wood Rails, White-tailed and Gartered Trogons and Citron-throated Toucan. We were kept busy overhead with Mealy and Orange-winged Parrots and both Blue-and- yellow and Chestnut-fronted Macaws. We teased out a male Black Antshrike, enjoyed a confiding Chestnut-backed Antbird and had great views of the localised Black-billed Flycatcher. Other species noted during what was a very busy day, included Rufous Mourner, Black-bellied Wren, Fulvous-vented Euphonia, Golden-hooded and Yellow-backed Tanagers, Golden-headed Manakin, Swallow-tailed Kite and King Vulture. Beautiful Woodpecker eluded us for much of the morning but then we located some calling right above the lodge and they soon showed well, while we also located White-fronted Nunbird and Cinnamon Woodpecker near one of the cabins. In the evening, we managed to tape in a couple of spectacular Spectacled Owls, while mammals included Humboldt's White-faced Capuchin, Central American Agouti and Brown Spider Monkey.

The following morning, we hiked out, after some early morning birding and then made the long drive to the Piha Reserve. On the walk out we managed an eleventh-hour Sooty Ant Tanager and also picked up some nice(Lesser) Pied Puffbirds. Much of the rest of the day was spent with the long drive north to the Piha reserve, though a short birding stop near the reserve did produce a group of White-footed Tamarins.

The next day we first drove to a ridge, where we enjoyed great views of the endemic Chestnut-capped Piha, before spending much of the day hiking the trails in the reserve. The trails produced a steady stream of interesting species including Western Woodhaunter, Spotted Woodcreeper, endemic Black-and-gold Tanager, stunning Purplish-mantled Tanager and a couple of Indigo Flowerpiercers. A Pale-vented Thrush was a real surprise and we also added White-crowned Manakin, Yellow-throated Chlorospingus, Red-headed Barbet, and Silver-throated Tanager. In the late afternoon we returned to the lodge, where we managed to see Purple-throated Woodstar, Greenish Puffleg and Crowned Woodnymph on the feeders.

Our final morning at the Piha reserve was spent birding both the road and the trails. The road birding was very productive, as we had a group of Blue-fronted Parrotlets fly through. We also got good looks at a group of three Scarlet-and-white Tanagers, Sooty Headed Wren and our first White-naped Brushfinches. A short walk on the trails added Green-fronted Lancebill, Ochre-breasted Antpitta and White-crowned Tapaculo. After an early lunch we slowly birded down the road below the lodge. A very obliging Magdalena Antbird was a real highlight and we also found four endemic White-mantled Barbets and the near-endemic Bar-crested Antshrike. Other species seen included Checker-throated Stipplethroat, White-bibbed and Striolated Manakins, Slaty Spinetail, Olivaceous Flatbill, Dusky-faced, White-shouldered, Tawny-crested and Guira Tanagers and Bay Wren. We then had a slight change of plan and rather than driving to our original destination we headed off to Santa Rosa de Osos on the trail of a completely new species.

The following morning, we met a local guide who whisked us off to a brand-new reserve. This reserve was supposed to hold our main target, the endemic and recently rediscovered Antioquia Brushfinch but was also said to host a new species of Antpitta. Shortly after arrival we were taken to a feeding station, where a very obliging individual of this new Antpitta showed well. It was clearly a Tawny Antpitta type but was more golden in colour, had a larger bill and was occurring in scrubby subparamo, rather than true paramo habitat. After taking our fill of this exciting new species, we spent more time looking for the Antioquia Brushfinch, which was rather elusive but

which eventually showed well. We also logged the local Black-throated Flowerpiercers, of the highly isolated *villeumieri* race, which is almost certainly a split in waiting. Other species seen included Flammulated Treehunter, Black-collared Jay, Grey-browed Brushfinch, Collared Inca and Golden-fronted Redstart (of the golden-fronted chrysops form, which is a potential split). Following this we began the drive to Urrao, which due to various protest and road problems, became extremely long and tortuous and we arrived at our Hotel late in the night.

The following morning, we still had to rise early, as we needed to head up into the mountain for the day. We began the ascent to the Colibri del Sol reserve, with some on horseback and some on foot. Birding opportunities during the climb were few and we headed straight for the reserve house, where a hearty breakfast awaited us. We watched the hummingbird feeders, while feasting on Arepas and hot chocolate and these held Long-tailed Sylph, Mountain Velvetbreast and Sword-billed Hummingbird. Most of our main targets were higher still, so we continued on up the increasingly steep slopes. Whether on horse or foot, I think that we were all happy to reach the upper feeders. Here a female of the endemic and highly localised Dusky Starfrontlet showed well and then; after continuing a bit further, we managed good views of the endemic Paramillo Tapaculo. A fine male Chestnut-bellied Flowerpiercer completed the trio of key endemics from this higher elevation and it was soon time to begin the much easier descent to the Lodge. Our final target was the endemic and highly localised Urrao Antpitta, which had been erratically visiting a feeding station. We had looked in the morning with no success but when we arrived, we were amazed to find it sitting there waiting to be fed. It had been a long day but with all four of the key endemics in the bag, it was a happy, if tired group, which began the final descent to Urrao. We added little else as we hiked out, though better views of Black-collared Jay and a fly through by some Rusty-faced Parrots, helped break up the hike.

The next day we left Urrao early and headed for the Cauca valley, where we hoped to mop up some Cauca Valley endemics. We quickly located Antioquia Wren and followed this with a couple of Greyish Piculets. Having found these key species so quickly, we had some time for general birding and commoner species seen included, Red-rumped Woodpecker, abundant Scarlet-fronted Parakeets, a couple of Cinereous Becards, a nice Slate-headed Tody-Flycatcher, a couple of Northern Mouse-coloured Tyrannulets, some Clay-coloured Thrushes and a couple of Black-striped Sparrows. We then continued on to Medellin, with a lunch stop adding Acorn Woodpecker, both Olive-grey and Black-winged Saltators and Flame-rumped Tanager. In the late afternoon we visited La Romera, where we quite quickly located an endemic male Yellow-headed Manakin and a few commoner species, including Andean Motmot.

The following morning most of the group opted to stay in the Hotel but one stalwart was keen for another visit to La Romera. This proved to be productive, with good views of the endemic Red-bellied Grackle, a lovely singing Andean Solitaire, abundant Sickle-winged Guans, Common Chlorospingus, Russet-crowned Warbler, Azara's Spinetail and Black-capped Tanager, among a host of commoner species. We then caught an early afternoon flight to Bogota and said goodbye to two of the group, who were connecting with international flights, while the rest of us retired to a Hotel for the night.

The next morning, we had a very early start to catch a very early flight to Mitu. This meant though, that we could have much of a day birding in Mitu. By mid-morning we were already watching a group of Maroon-tailed Parakeets by the side of the road, on the outskirts of the town. We then birded around a covered bridge, where we quickly found Yellow-bellied Dacnis, Amazonian Scrub Flycatcher, Bronzy Jacamar and Cherries Antwren. Swallow-winged Puffbirds hawked from most available perches, Short-tailed and Fork-tailed Palm Swifts passed overhead, our first Greater Yellow-headed Vultures cruised by and Black-chinned Antbirds skulked along the waterways. We returned to the same area in the afternoon and this time ventured further into the community. A female Black-bellied Thorntail was a bonus as was a Dusky-billed Parrotlet, which was mixed with a flock of Cobalt-winged Parakeets, both being generally scarce or possibly overlooked in the Mitu area. We also found Gilded Barbet, Black Caracara, Black-headed Parrots, lots of Red-bellied Macaws and a pair of scarlet Macaws. A male Spangled Cotinga perched on a distant tree and a young and quite confident White-browed Purplethroat was much appreciated. We logged a veritable avalanche of new flycatchers with Yellow-browed Tody-Flycatcher, White-lored Tyrannulet, Slender-footed Tyrannulet and Amazonian Tyrannulet. A stealthy group of Azure-naped Jays were a key target and a species which can often be tricky to locate. The same can be said for a singing Plumbeous Euphonia, while other Euphonias seen included Golden-bellied, Rufous-bellied and White-lored Euphonia. New Tanagers featured with Masked, Turquoise and Opal-rumped all seen well, while we also recorded both Chestnut-bellied Seed Finch and Chestnut-bellied Seedeater. After a couple of weeks in the Andes, to suddenly be presented with so many new species in such a short time was quite exhilarating!

The following day we headed out into the infamous and sometimes unforgiving Mitu Cachivera trail. We left especially early and this proved to be very beneficial, as soon after dawn we located one of the key Mitu targets, a vocal and surprisingly responsive Bar-bellied Woodcreeper. We followed this with a couple of Grey-bellied Antbirds that skulked at our feet and then had brief views of one of the holy

grails, a male Purple-breasted Cotinga, which perched out for a short period. The tone was set for the rest of the morning and the targets kept coming thick and fast. Black Manakin showed well, Citron-bellied Attila performed admirably and we enjoyed good views of both Yellow-crested Manakin and Saffron-crowned Tyrant-Manakins. A pair of fairly responsive Chestnut-crested Antbirds also made several passes. This species can be notoriously difficult to locate, so to see them on our first full day and at an unexpected site, was especially fortuitous. We also found Lettered and Many-banded Aracaris, Amazonian Antshrike, Rufous-backed Stipplethroat, Spot-backed Antwren, Black-throated Antbird, Blue-capped Manakin and an obliging female Pompadour Cotinga. Flycatchers included Fuscous (the *duidae* race), Dusky-chested, Sulphury and Yellow-throated, while we also added White-eyed Tody-tyrant, Brown-headed Greenlet and Fulvous-crested and Paradise Tanagers. It really was a superlative mornings birding, with a surfeit of key targets logged on a trail, which can at times be very hard birding. In the afternoon we visited the Bocotoma trail where birding was inevitably slower. The only significant additions were a Brown-winged Schiffornis and a brief White-chinned Woodcreeper, although we did enjoy good views of a few of the species that we had seen earlier.

The following morning, we left very early and headed to a well-known bridge. We waited in the half light and after a short time enjoyed great views of a male Fiery Topaz, as it danced over the water. Other species seen in the area included Reddish Hermit, Black-eared fairy and then a real surprise, with an Orinoco Piculet, located feeding at the forest edge. We also spent some time on a skulking Black Bushbird, which eventually showed well, representing our second Bushbird species of the trip! Olive Oropendolas were seen flying over and Amazonian Grosbeaks sulked in the understorey. We then moved on to the Pueblo Nuevo area, where we birded several trails in the late morning and again in the afternoon. We found a multitude of species including Chestnut-eared Aracari, Channel-billed and White-throated Toucans, Red-fan Parrot, Imeri Warbling Antbird, a skulking male Black-headed Antbird and a Short-billed Leaf-tosser. Mixed flocks held Olive-Backed Foliage-gleaner and Eastern Woodhaunter and we also picked up Magpie Tanager, Red-throated Caracara and White-crowned Manakin.

The following day we drove a long way out of Mitu and started birding along a remote road surrounded by primary forest. This produced Paradise Jacamar, White-fronted Nunbird, Ivory-billed Aracari, Red-necked, Waved and Chestnut Woodpeckers and Pink-throated Becard. A vocal pair of Ochre-throated Foliage-gleaners showed well and we finally caught up with Grey Antbird, while migrant Olive-sided Flycatchers mixed with the Jacamars on the overhead power lines. We then birded a nearby trail, where we added Great Jacamar, Rufous-tailed Xenops and Cinnamon-rumped Foliage-gleaner. A key target was Yellow-throated Antwren and after much neck straining, we were able to observe this canopy loving species. Lunch by a hydro-electric scheme gave us a brief respite from the forest birding and the impressive rapids held Black-collared, White-banded and White-winged Swallow, Large-billed Tern and Pied Plover. The afternoon was inevitable much quieter but did produce a major target with a male Tawny-tufted Toucanet eventually showing well. Mammals seen during the day included some quite confiding Lucifer Titis and a Common Opossum, which was foraging along the roadside early in the morning.

The following day we returned to the Pueblo Nuevo area where we first visited a favoured Parrot feeding area. Here we saw hundreds of Orange-cheeked Amazons while a couple of Kewell's Parrots flew over. We also picked up the much-desired Green-tailed Goldenthrout and added (Greater) Pied Puffbird and Speckled Chachalaca. The rest of the day was spent along various trails in the Pueblo Nuevo area. These were rather quiet at times but we did manage to eke out White-chested Puffbird, Pearly Antshrike, Rufous-capped Antthrush and a *Duidae* Woodcreeper.

Our last morning in Mitu was spent on the Bocotoma Trail, where we had good views of some skulking Collared Gnatwrens and a lovely Ringed Antpipit. We also found Buff-throated Woodcreeper and Rufous-tailed Flatbill, while a quick stop on the drive back to the Hotel produced Point-tailed Palmcreeper. We then caught a flight to Bogota, where upon arrival we had to immediately set off down the eastern Andes. Road closures meant that we would not be able to access the area that we wanted to visit the next day, unless we spent the night nearby. After battling various traffic problems, we eventually made a late arrival at our Hotel. It was not ideal but it would mean that we could chase another endemic the following day.

Our final morning involved a very steep jeep ride high into the mountains, where we visited a Finca that had a hand fed endemic Cundinamarca Antpitta. The Antpitta performed admirably shortly after our arrival and we also logged the localised Ochre-breasted Brushfinch in the garden. We had time to bird some of the trails and this produced a brief Muisca Antpitta and a couple of Long-tailed Tapaculos. A pair of Rufous-banded Owls were also seen at roost and we added White-throated Toucanet and nice views of more Inca and Black-collared Jays. It was then time to leave, allowing plenty of time for the climb back up to Bogota, where our wonderful Colombia adventure finished in the afternoon. Despite various hiccups, we had managed to visit many remote regions of Colombia and to see

some of its lesser-known endemics and specialities. It really had been an intoxicating cocktail of quality birds and fantastic scenery, a truly memorable adventure.

SYSTEMATIC LIST OF SPECIES RECORDED DURING THE TOUR

Species marked with the diamond symbol (◊) are either endemic to the country or local region or considered 'special' birds for some other reason (e.g., it is only seen on one or two Birdquest tours; it is difficult to see across all or most of its range; the local form is endemic or restricted-range and may in future be treated as a full species).

The species names and taxonomy used in the bird list follows Gill, F., Donsker, D., & Rasmussen, P.(Eds). **2024. IOC World Bird List (v14.1).**

Where the subspecies seen is/are known, these are often given in parentheses at the end of the species comment.

Species only likely on the Cundinamarca Antpitta & Mitú post-tour Extension are marked with the symbol E.

BIRDS

Great Tinamou *Tinamus major* Heard.

White-throated Tinamou *Tinamus guttaus* Heard.

Highland Tinamou ◊ *Nothocercus bonapartei* A single seen on the trail at the Cerulean Warbler Reserve.

Tawny-breasted Tinamou ◊ *Nothocercus julius* Heard.

Cinereous Tinamou *Crypturellus cinereus* Heard.

Little Tinamou *Crypturellus soui* Heard.

Grey-legged Tinamou ◊ *Crypturellus duidae* Heard.

Variegated Tinamou *Crypturellus variegatus* Heard.

Northern Screamer ◊ *Chauna chavaria* Seen on the drives in and out of Paujil, maximum five.

Black-bellied Whistling Duck *Dendrocygna autumnalis* Seen at a lunch stop in the Magdalena Valley.

Speckled Chachalaca *Ortalis guttata* Seen at the Parrot stakeout at Mitu.

Colombian Chachalaca ◊ *Ortalis columbiana* Endemic. Seen on multiple days during the trip.

Band-tailed Guan ◊ *Penelope argyrotis* Four were seen at dusk at the Bushbird Reserve.

Andean Guan ◊ *Penelope montagnii* A couple were seen on our first day in the Perija.

Spix's Guan *Penelope jacquacu*

Wattled Guan ◊ *Aburria aburri* Heard.

Sickle-winged Guan ◊ *Chamaepetes goudotii* Seen commonly at multiple sites.

Blue-billed Curassow ◊ *Crax alberti* Endemic. A minimum of twelve at were seen at Paujil.

Crested Bobwhite ◊ *Colinus cristatus* Four were seen well at Los Besotes.

Marbled Wood Quail *Odontophorus gujanensis* Heard below the lodge at the Piha Reserve.

Black-fronted Wood Quail ♦ *Odontophorus atrifrons* Non leader. One group member stayed behind at the Perija lodge on one afternoon and saw two visiting the feeder.

Chestnut Wood Quail ♦ *Odontophorus hyperythrus* Endemic. Heard.

Gorgeted Wood Quail ♦ *Odontophorus strophium* Endemic. Heard.

Short-tailed Nighthawk *Lurocalis semitorquatus* Seen at Mitu.

Rufous-bellied Nighthawk *Lurocalis rufiventris* A single perched on fence posts at the Cerulean Warbler Reserve.

Blackish Nightjar *Nyctipolus nigrescens* Seen at Mitu.

Pauraque *Nyctidromus albicollis*

Band-winged Nightjar *Systellura longirostris* Seen at several sites, with best views in the Perija.

Lyre-tailed Nightjar ♦ *Uropsalis lyra* A female briefly at dawn at the Piha Reserve.

Rufous Nightjar *Antrostomus rufus* Heard.

Common Potoo *Nyctibius griseus* Heard.

White-collared Swift *Streptoprocne zonaris*

Grey-rumped Swift *Chaetura cinereiventris*

Short-tailed Swift *Chaetura brachyura*

White-tipped Swift *Aeronautes montivagus* Seen at several Andean sites.

Fork-tailed Palm Swift *Tachornis squamata*

Fiery Topaz ♦ *Topaza pyra* A dazzling male at Mitu.

White-necked Jacobin *Florisuga mellivora*

Stripe-throated Hermit *Phaethornis striigularis*

Reddish Hermit *Phaethornis ruber*

Sooty-capped Hermit *Phaethornis augusti*

Pale-bellied Hermit *Phaethornis anthophilus* Heard.

Green Hermit *Phaethornis guy*

Great-billed Hermit *Phaethornis malaris*

Green-fronted Lancebill ♦ *Doryfera ludovicae* A single showed well at the Piha Reserve.

Brown Violetear *Colibri delphinae*

Lesser Violetear *Colibri cyanotus*

Sparkling Violetear *Colibri coruscans*

Black-eared Fairy *Heliophryx auritus*

Green-tailed Goldenthrroat ♦ *Polytmus theresiae* A single at the Parrot stakeout at Mitu.

Black-throated Mango *Anthracothorax nigricollis*

Longuemare's Sunangel ♦ *Helianthus clarisse* Common at Soata and in the Perija.

Tourmaline Sunangel *Helianthus exortis*

Black-bellied Thorntail ♦ *Discosura langsdorffi* A couple of females on separate days at Mitu were a tour highlight.

Speckled Hummingbird *Adelomyia melanogenys*

Long-tailed Sylph *Agelaiocercus kingii*

Tyrian Metaltail *Metallura tyrianthina* The nominate race was seen in the Andes, while the *districta* form was seen in the Perija – a potential split.

Perija Metaltail ♦ *Metallura iracunda* Multiple individuals were seen on the Paramo in the Perija.

Greenish Puffleg ♦ *Haplophedya aureliae* Seen at the Piha Reserve and La Romera.

Black Inca ♦ *Coeligena prunellei* Endemic. Seen on the feeders at the Cerulean Warbler Reserve.

Collared Inca *Coeligena torquata*

Dusky Starfrontlet ♦ *Coeligena orina* Endemic. A couple showed well after our long hike at the Colibri del Sol Reserve.

Perija Starfrontlet ♦ *Coeligena consita* A minimum of three, including a dazzling male, at the Perija Reserve, were a trip highlight.

Mountain Velvetbreast *Lafresnaya lafresnayi*

Sword-billed Hummingbird *Ensifera ensifera* This iconic species showed well at Colibri Del Sol.

Buff-tailed Coronet *Boissonneaua flavescens*

White-booted Racket-tail *Ocreatus underwoodii*

Fawn-breasted Brilliant ♦ *Heliodoxa rubinoides*

Green-crowned Brilliant *Heliodoxa jacula*

Purple-throated Woodstar ♦ *Philodice mitchellii*

White-bellied Woodstar *Chaetocercus mulsant*

Rufous-shafted Woodstar ♦ *Chaetocercus jourdanii* Several were seen visiting flowering bushes in the Perija.

Red-billed Emerald ♦ *Chlorostilbon gibsoni*

Coppery Emerald ♦ *Chlorostilbon russatus* A single was seen on the descent from the Perija Lodge.

Short-tailed Emerald ♦ *Chlorostilbon poortmani* A single was seen at the Cerulean Reserve.

White-vented Plumeleteer *Chalybura buffonii*

Crowned Woodnymph ♦ (Violet-c W) *Thalurania [colombica] colombica*

Crowned Woodnymph ♦ (Green-c W) *Thalurania [colombica] fannyae*

Fork-tailed Woodnymph *Thalurania furcata*

Steely-vented Hummingbird *Saucerottia saucerottei*

Indigo-capped Hummingbird ♦ *Saucerottia cyanifrons* Endemic. Seen in the Chicamocha canyon and at the Cerulean Reserve.

Chestnut-bellied Hummingbird ♦ *Saucerottia castaneiventris* Endemic. Abundant at the feeders in the Chicamocha canyon.

Rufous-tailed Hummingbird *Amazilia tzacatl*

Andean Emerald *Uranomitra franciae*

Shining-green Hummingbird ♦ *Chrysuronia goudoti* Seen on both visits to Los Besotes Reserve.

Versicolored Emerald *Chrysuronia versicolor*

Blue-chested Hummingbird *Polyerata amabilis*

Violet-bellied Hummingbird *Chlorestes julie*

Greater Ani *Crotophaga major*

Smooth-billed Ani *Crotophaga ani*

Striped Cuckoo *Tapera naevia*

Pavonine Cuckoo *Dromococcyx pavoninus* Heard.

Squirrel Cuckoo *Piaya cayana*

Black-bellied Cuckoo *Piaya melanogaster*

Rock Dove (introduced) *Columba livia*

Band-tailed Pigeon *Patagioenas fasciata*

Pale-vented Pigeon *Patagioenas cayennensis*

Plumbeous Pigeon *Patagioenas plúmbea* Non leader.

Ruddy Pigeon *Patagioenas subvinacea*

Scaled Dove *Columbina squammata*

Common Ground Dove *Columbina passerina*

Plain-breasted Ground Dove *Columbina minuta*

Ruddy Ground Dove *Columbina talpacoti*

Blue Ground Dove *Claravis pretiosa*

Maroon-chested Ground Dove ♦ *Paraclaravis mondetoura* Heard.

Ruddy Quail-Dove *Geotrygon montana* A single was seen well at the Cerulean Reserve.

White-tipped Dove *Leptotila verreauxi*

Grey-fronted Dove *Leptotila rufaxilla* Heard.

Lined Quail-Dove ♦ *Zentrygon linearis* Heard.

Eared Dove *Zenaida auriculata*

Grey-cowled Wood Rail *Aramides cajaneus*

Purple Gallinule *Porphyrio martinica*

Double-striped Thick-knee *Burhinus bistriatus* A couple of the Los Besotes entrance track

Southern Lapwing *Vanellus chilensis*

Pied Plover (P Lapwing) *Hoploxypterus cayanus* A couple at Mitu.

Wattled Jacana *Jacana jacana*

Solitary Sandpiper *Tringa solitaria*

Neotropic Cormorant *Nannopterum brasilianum*

Green Ibis *Mesembrinibis cayennensis* A single on the drive into Paujil.

Bare-faced Ibis *Phimosus infuscatus*

Rufescent Tiger Heron *Tigrisoma lineatum* A single on the drive into Paujil.

Striated Heron *Butorides striata*

Western Cattle Egret *Bubulcus ibis*

Cocoi Heron *Ardea cocoi*

Great Egret (American G E) *Ardea [alba] egretta*

Little Blue Heron *Egretta caerulea*

Snowy Egret *Egretta thula*

King Vulture *Sarcoramphus papa* Seen at Paujil and Mitu.

Andean Condor ♦ *vulture gryphus* Seen in the Perija mountains.

Black Vulture *Coragyps atratus*

Turkey Vulture *Cathartes aura*

Lesser Yellow-headed Vulture *Cathartes burrovianus*

Greater Yellow-headed Vulture *Cathartes melambrotus*

Osprey *Pandion haliaetus*

White-tailed Kite *Elanus leucurus*

Grey-headed Kite *Leptodon cayanensis*

Swallow-tailed Kite *Elanoides forficatus*

Black Hawk-Eagle *Spizaetus tyrannus* Seen overhead at Paujil and the Cerulean Reserve.

Black-and-chestnut Eagle *Spizaetus isidori* A brief single in the Perija mountains.

Double-toothed Kite *Harpagus bidentatus*

Plain-breasted Hawk *Accipiter ventralis*

Plumbeous Kite *Ictinia plumbea*

Savanna Hawk *Buteogallus meridionalis*

Roadside Hawk *Rupornis magnirostris*

Black-chested Buzzard-Eagle *Geranoaetus melanoleucus* A couple over the Paramo in the Perija.

Grey-lined Hawk *Buteo nitidus*

Broad-winged Hawk *Buteo platypterus*

Short-tailed Hawk *Buteo brachyurus*

Andean Pygmy Owl *Glaucidium jadinii* A single showed well on the Paramo in the Perija.

Ferruginous Pygmy Owl *Glaucidium brasilianum* Heard at Los Besotes.

White-throated Screech Owl *Megascops albogularis* A pair were seen well at Soata.

Tropical Screech Owl *Megascops choliba*

Cinnamon Screech Owl ♦ *Megascops petersoni* Seen well at the Cerulean Reserve.

Choco Screech Owl ♦ *Megascops centralis* Seen well at Paujil.

Spectacled Owl *Pulsatrix perspicillata* A pair were seen well at Paujil.

Mottled Owl *Strix virgate* Seen well at the Cerulean Reserve.

Black-and-white Owl ♦ *Strix nigrolineata* A single on a day roost at the Cerulean Reserve.

Rufous-banded Owl ♦ *Strix albitarsis* Non-leader. A roosting pair were shown to the group by the local guide at Finca La Herrea.

Pavonine Quetzal *Pharomachrus pavoninus* Heard.

Golden-headed Quetzal *Pharomachrus auriceps* Heard.

White-tailed Trogon (Western W-t T) *Trogon chionurus*

Green-backed Trogon *Trogon viridis* Heard.

Gartered Trogon (Northern Violaceous T) *Trogon caligatus*

Amazonian Trogon *Trogon ramonianus* Heard.

Collared Trogon *Trogon collaris*

Masked Trogon *Trogon personatus*

Amazon Kingfisher *Chloroceryle amazona*

Green Kingfisher *Chloroceryle americana*

Ringed Kingfisher *Megaceryle torquata* Non leader.

Andean Motmot ♦ (Highland M) *Momotus aequatorialis*

Rufous Motmot *Baryphthengus martii* Non leader.

Rufous-tailed Jacamar *Galbula ruficauda* Heard.

Bronzy Jacamar ♦ *Galbula leucogastra* Seen well at Mitu.

Paradise Jacamar *Galbula dea* Seen well at Mitu

Great Jacamar *Jacamerops aureus* Seen well at Mitu

Pied Puffbird *Notharchus tectus* 'Lesser' Pied Puffbirds were seen on the drive out of Paujil and a 'Greater' Pied Puffbird was seen at Mitu.

Barred Puffbird ♦ *Nystalus radiates* A single seen well in the Cacao plantations below San Vicente.

Russet-throated Puffbird ◊ *Hypnelus ruficollis* Heard

White-chested Puffbird ◊ *Malacoptila fusca* A single seen well at Mitu.

Moustached Puffbird ◊ *Malacoptila mystacalis* Seen well on multiple days of the drip, including in the Perija and at the Bushbird Reserve.

White-fronted Nunbird *Monasa morphoeus* The sclateri subspecies was seen at Paujil and the peruana form was seen at Mitu. A cis-Andean split is possible for this species in the future.

Swallow-winged Puffbird *Chelidoptera tenebrosa*

White-mantled Barbet ◊ *Capito hypoleucus* Four were seen below the Piha Reserve.

Gilded Barbet *Capito auratus*

Red-headed Barbet *Eubucco bourcierii*

White-throated Toucanet ◊ (Andean T) *Aulacorhynchus [albivitta] albivitta* Heard.

Crimson-rumped Toucanet ◊ *Aulacorhynchus haematopygus* Heard.

Lettered Aracari *Pteroglossus inscriptus*

Ivory-billed Aracari *Pteroglossus azara*

Chestnut-eared Aracari *Pteroglossus castanotis*

Many-banded Aracari *Pteroglossus pluricinctus*

Collared Aracari *Pteroglossus torquatus*

Tawny-tufted Toucanet *Selenidera nattereri* A rather shy individual was eventually seen well at Mitu.

Channel-billed Toucan *Ramphastos vitellinus*

Citron-throated Toucan ◊ *Ramphastos citreolaemus*

Keel-billed Toucan *Ramphastos sulfuratus* Heard.

White-throated Toucan *Ramphastos tucanus*

Yellow-throated Toucan (Chestnut-mandibled T) *Ramphastos ambiguous* Heard.

Orinoco Piculet ◊ *Picumnus pumilus* A single seen well at the Fiery Topaz bridge at Mitu.

Olivaceous Piculet *Picumnus olivaceus*

Greyish Piculet ◊ *Picumnus granadensis* Endemic. A couple were seen in the Cauca valley.

Chestnut Piculet ◊ *Picumnus cinnamomeus* Non Leader

Acorn Woodpecker *Melanerpes formicivorus*

Yellow-tufted Woodpecker *Melanerpes cruentatus*

Beautiful Woodpecker ◊ *Melanerpes pulcher* Endemic. Seen well at Paujil

Red-crowned Woodpecker *Melanerpes rubricapillus*

Red-rumped Woodpecker *Veniliornis kirkii*

Red-stained Woodpecker *Veniliornis affinis*

Smoky-brown Woodpecker *Leuconotopicus fumigatus*

Yellow-throated Woodpecker *Piculus flavigula*

Golden-olive Woodpecker *Colaptes rubiginosus*

Crimson-mantled Woodpecker *Colaptes rivolii*

Spot-breasted Woodpecker *Colaptes punctigula*

Cinnamon Woodpecker *Celeus loricatus* A couple were seen at Paujil.

Waved Woodpecker *Celeus [undatus] grammicus*

Chestnut Woodpecker *Celeus elegans*

Lineated Woodpecker *Dryocopus lineatus*

Powerful Woodpecker *Campephilus pollens* A pair were seen in the oak forests above Soata.

Red-necked Woodpecker *Campephilus rubricollis*

Crimson-crested Woodpecker *Campephilus melanoleucos* Non leader.

Black Caracara *Daptrius ater*

Red-throated Caracara *Ibycter americanus*

Crested Caracara (Northern C C) *Caracara [plancus] cheriway*

Yellow-headed Caracara *Milvago chimachima*

Laughing Falcon *Herpetotheres cachinnans*

Collared Forest Falcon *Micrastur semitorquatus* Heard.

American Kestrel *Falco sparverius*

Blue-fronted Parrotlet ♦ *Touit dilectissimus* Heard at Cerulean and a group seen in flight at the Piha reserve.

Barred Parakeet *Bolborhynchus lineola* Heard.

Orange-chinned Parakeet *Brotogeris jugularis*

Cobalt-winged Parakeet *Brotogeris cyanoptera*

Orange-cheeked Parrot ♦ *Pyrilia barrabandi* Impressive number seen at Mitu.

Rusty-faced Parrot ♦ *Hapalopsittaca amazonina* Small flocks seen at Soata and Colibri del Sol.

White-capped Parrot *Pionus seniloides* Heard.

Blue-headed Parrot *Pionus menstruus*

Bronze-winged Parrot *Pionus chalcopterus*

Scaly-naped Amazon *Amazona mercenarius*

Mealy Amazon *Amazona farinose*

Kawall's Amazon ♦ *Amazona kawalli* Two were seen in flight at Mitu.

Orange-winged Amazon *Amazona amazonica*

Dusky-billed Parrotlet *Forpus modestus* Seen on a couple of days at Mitu.

Spectacled Parrotlet *Forpus conspicillatus*

Black-headed Parrot *Pionites melanocephalus*

Red-fan Parrot ◊ *Deroptyus accipitrinus*

Painted Parakeet ◊ (Todd's P) *Pyrrhura [picta] subandina* A very confiding flock near La Jagua.

Maroon-tailed Parakeet *Pyrrhura melanura*

Brown-throated Parakeet *Eupsittula pertinax*

Red-bellied Macaw *Orthopsittaca manilatus*

Blue-and-yellow Macaw *Ara ararauna*

Chestnut-fronted Macaw *Ara severus*

Military Macaw *Ara militaris* A small flock in flight at the base of the Perija.

Scarlet Macaw *Ara macao*

Scarlet-fronted Parakeet *Psittacara wagleri*

Short-billed Leaf-tosser *Sclerurus rufularis* Seen well at Mitu.

Olivaceous Woodcreeper *Sittasomus griseicapillus*

Tyrannine Woodcreeper *Dendrocincla tyrannina*

White-chinned Woodcreeper *Dendrocincla merula*

Plain-brown Woodcreeper *Dendrocincla fuliginosa*

Wedge-billed Woodcreeper *Glyphorhynchus spirurus*

Black-banded Woodcreeper *Dendrocolaptes picumnus* Heard.

Bar-bellied Woodcreeper ◊ *Hylexetastes stresemanni* Seen very well at Mitu, a trip highlight.

Strong-billed Woodcreeper *Xiphocolaptes promeropirhynchus* Heard.

Buff-throated Woodcreeper *Xiphorhynchus guttatus*

Cocoa Woodcreeper *Xiphorhynchus susurrans*

Black-striped Woodcreeper *Xiphorhynchus lachrymosus* Heard.

Spotted Woodcreeper *Xiphorhynchus erythropygius*

Olive-backed Woodcreeper *Xiphorhynchus triangularis*

Straight-billed Woodcreeper *Dendroplex picus*

Brown-billed Scythebill *Campylorhamphus pusillus* Seen at Cerulean and the Piha Reserve.

Streak-headed Woodcreeper *Lepidocolaptes souleyetii*

Montane Woodcreeper *Lepidocolaptes lacrymiger*

Duida Woodcreeper *Lepidocolaptes duidae*

Slender-billed Xenops *Xenops tenuirostris*

Plain Xenops *Xenops genibarbis*

Streaked Xenops *Xenops rutilans*

Point-tailed Palmcreeper *Berlepschia rikeri* Seen on the outskirts of Mitu.

Rufous-tailed Xenops *Microxenops milleri*

Streaked Tuftedcheek *Pseudocolaptes boissonneautii*

Rusty-winged Barbtail ◊ *Premnornis guttuliger* Heard.

Caribbean Hornero ◊ *Furnarius longirostris*

Cinnamon-rumped Foliage-gleaner *Philydor pyrrhodes* Seen well at Mitu.

Montane Foliage-gleaner *Anabacerthia striaticollis*

Lineated Foliage-gleaner *Syndactyla subalaris*

Buff-fronted Foliage-gleaner *Dendroma rufa*

Flammulated Treehunter *Thripadectes flammulatus* Seen well at the 'new' Antpitta Reserve.

Striped Treehunter ◊ *Thripadectes holostictus* Heard.

Streak-capped Treehunter ◊ *Thripadectes virgaticeps*

Ochre-throated Foliage-gleaner* *Automolus ochrolaemus* Seen well at Mitu.

Eastern Woodhaunter *Automolus subulatus* Seen well at Mitu

Western Woodhaunter *Automolus virgatus* Seen well at the Piha Reserve.

Olive-backed Foliage-gleaner *Automolus infuscatus* Seen at Mitu.

Spotted Barbtail *Premnoplex brunnescens* Heard.

Pearled Treerunner *Margarornis squamiger*

White-browed Spinetail ◊ *Hellmayrea gularis*

Perija Thistletail ◊ *Asthenes perijana* Seen well on the Paramo above the Perija.

Double-banded Greytail ◊ *Xenerpestes minlosi* Seen well below the Cerulean Reserve.

Slaty Spinetail *Synallaxis brachyura*

Pale-breasted Spinetail *Synallaxis albescens* Heard.

Azara's Spinetail *Synallaxis azarae*

White-whiskered Spinetail ◊ *Synallaxis candei* Seen well at Los Besotes.

Rufous Spinetail *Synallaxis unirufa* The munoztebari subspecies that we saw in the Perija is vocally distinct and a probable split.

Stripe-breasted Spinetail ◊ *Synallaxis cinnamomea* Seen well at the Bushbird Reserve.

Rufous-rumped Antwren ◊ *Euchrepomis callinota* Seen at the Cerulean Reserve.

Black Bushbird *Neotantes niger* Seen well at Mitu.

Recurve-billed Bushbird ◊ *Clytoctantes alixii* Seen well at the Bushbird Reserve.

Checker-throated Stipplethroat *Epinecrophylla fulviventris*

Rufous-backed Stipplethroat ◊ *Epinecrophylla [haematonota] pyrrhonota*

Grey-bellied Antbird ◊ *Ammonastes pelzelni* A pair showed well at Mitu.

Black-throated Antbird *Myrmophylax atrothorax*

Pygmy Antwren *Myrmotherula brachyuran* Non Leader.

Pacific Antwren ◊ (P Streaked A) *Myrmotherula pacifica*

Cherrie's Antwren ◊ *Myrmotherula cherriei* Seen well at Mitu.

Yellow-throated Antwren ◊ *Myrmotherula ambigua* Seen in the canopy at Mitu

White-flanked Antwren *Myrmotherula axillaris* Heard.

Slaty Antwren *Myrmotherula schisticolor*

Northern White-fringed Antwren *Formicivora intermedia*

Plain-throated Antwren *Isleria hauxwelli* Heard.

Dusky-throated Antshrike *Thamnomanes ardesiacus* Heard.

Cinereous Antshrike *Thamnomanes caesius* Heard.

Pearly Antshrike *Megastictus margaritatus* Seen well at Mitu.

Spot-backed Antwren *Herpsilochmus dorsimaculatus* Seen well at Mitu.

Plain Antvireo *Dysithamnus mentalis*

Black-backed Antshrike ◊ *Thamnophilus melanonotus* Seen well at Los Besotes.

Bar-crested Antshrike ◊ *Thamnophilus multistriatus*

Black Antshrike ◊ *Thamnophilus nigriceps* Seen well at Paujil.

Uniform Antshrike *Thamnophilus unicolor*

Mouse-colored Antshrike *Thamnophilus murinus* Non leader.

Black-crowned Antshrike (Western Slaty A) *Thamnophilus atrinucha*

Amazonian Antshrike *Thamnophilus amazonicus*

Black-crested Antshrike *Sakesphorus canadensis*

Fasciated Antshrike *Cymbilaimus lineatus* Heard.

Great Antshrike *Taraba major* Heard.

Bicolored Antbird *Gymnopithys bicolor* Non leader.

Chestnut-crested Antbird ◊ *Rhegmatorhina cristata* A pair were seen well at Mitu.

Common Scale-backed Antbird *Willisornis poecilinotus* Heard

Klages's Antbird ◊ *Drymophila klagesi* Heard in the Perija and seen well at the Bushbird Reserve.

Imeri Warbling Antbird ◊ *Hypocnemis flavescens*

Parker's Antbird ◊ *Cercomacroides parkeri* Endemic. Seen well at the Cerulean and Piha Reserves.

Dusky Antbird *Cercomacroides tyrannina* Heard.

Grey Antbird *Cercomacra cinerascens*

Jet Antbird *Cercomacra nigricans* A male showed well on the drive to the Cerulean Reserve.

Black-chinned Antbird *Hypocnemoides melanopogon*

Chestnut-backed Antbird *Poliocrania exsul*

Magdalena Antbird ◊ *Sipia palliate* Seen well below the Piha Reserve.

Black-faced Antbird *Myrmoborus myotherinus* Heard.

Bare-crowned Antbird ◊ *Gymnocichla nudiceps* Heard.

Black-headed Antbird ◊ *Percnostola rufifrons* Seen well at Mitu.

Blue-lored Antbird ◊ *Hafferia immaculate* A skulking pair at the Cerulean Reserve.

Rufous-capped Antthrush *Formicarius colma*

Chestnut-crowned Antpitta *Grallaria ruficapilla*

Cundinamarca Antpitta ◊ *Grallaria kaestneri* Endemic. Seen well at Finca La Herrera.

Chestnut-naped Antpitta *Grallaria nuchalis* Heard.

White-bellied Antpitta ◊ *Grallaria hypoleuca* Seen at the Cerulean Reserve.

Perija Antpitta ◊ *Grallaria saltuensis* Seen in the Paramo at the Perija.

Muisca Antpitta *Grallaria rufula* Seen at Finca La Herrera

Chami Antpitta ◊ *Grallaria alvarezi* Endemic. Heard.

'Paisa' Antpiits ◊ (*Sp novum*) Endemic. Great views of the undescribed species at a new reserve in NW Antioquia.

Urrao Antpitta ◊ *Grallaria urraoensis* Endemic. Seen well at Colibri del Sol.

Thrush-like Antpitta *Myrmothera campanisona* Heard.

Ochre-breasted Antpitta *Grallaricula flavirostris* Seen well at the Piha Reserve.

Slaty-crowned Antpitta *Grallaricula nana* Heard.

Chestnut-belted Gnateater *Conopophaga aurita* Heard.

Chestnut-crowned Gnateater ◊ *Conopophaga castaneiceps* Seen well at the Cerulean and Piha Reserves.

Paramillo Tapaculo ◊ *Scytalopus canus* Endemic. Seen well at Colibri del Sol.

White-crowned Tapaculo ◊ (Northern W-c T) *Scytalopus atratus* Seen at the Piha Reserve.

Long-tailed Tapaculo ◊ *Scytalopus micropterus* Heard at Cerulean and seen at Finca La Herrera.

Blackish Tapaculo ◊ *Scytalopus latrans* Heard.

Magdalena Tapaculo ◊ (Upper M T) *Scytalopus rodriguezi* Endemic. Seen at the Cerulean Warbler Reserve.

Stiles's Tapaculo ♦ *Scytalopus stilesi* Endemic. Heard.

Pale-bellied Tapaculo ♦ (Mattoral T) *Scytalopus griseicollis* Seen well at Soata.

Perija Tapaculo ♦ *Scytalopus perijanus* Seen well in the Perija.

Spillmann's Tapaculo *Scytalopus spillmanni* Heard.

Wing-barred Piprites (W-b Manakin) *Piprites chloris* Heard.

Sooty-headed Tyrannulet *Phyllomyias griseiceps*

Plumbeous-crowned Tyrannulet ♦ *Phyllomyias plumbeiceps* Seen at the Cerulean Reserve.

Black-capped Tyrannulet *Phyllomyias nigrocapillus*

Yellow-crowned Tyrannulet *Tyrannulus elatus*

Forest Elaenia *Myiopagis gaimardii*

Amazonian Elaenia *Myiopagis cinerea* Heard.

Greenish Elaenia *Myiopagis viridicata*

Yellow-bellied Elaenia *Elaenia flavogaster*

Mountain Elaenia *Elaenia frantzii*

Brown-capped Tyrannulet *Ornithion brunneicapillus* Heard.

White-lored Tyrannulet *Ornithion inermis*

Southern Beardless Tyrannulet *Camptostoma obsoletum*

White-throated Tyrannulet *Mecocerculus leucophrys*

Northern Mouse-colored Tyrannulet *Nesotriccus incomtus*

Ringed Antpipit *Corythopis torquatus* Seen well at Mitu.

Spectacled Tyrannulet ♦ *Zimmerius improbus* Seen commonly in the Perija.

Slender-footed Tyrannulet *Zimmerius gracilipes*

Golden-faced Tyrannulet *Zimmerius chrysops*

Coopmans's Tyrannulet ♦ *Zimmerius minimus*

Variegated Bristle Tyrant *Pogonotriccus poecilotis*

Marble-faced Bristle Tyrant *Pogonotriccus ophthalmicus*

Streak-necked Flycatcher *Mionectes striaticollis*

Olive-striped Flycatcher *Mionectes galbinus*

Ochre-bellied Flycatcher *Mionectes oleagineus*

Sepia-capped Flycatcher *Leptopogon amaurocephalus*

Slaty-capped Flycatcher *Leptopogon superciliaris*

Slender-billed Inezia ♦ *Inezia tenuirostris* Seen at Los Besotes.

Amazonian Inezia *Inezia subflava*

Pale-tipped Inezia ♦ *Inezia caudata* Seen at Los Besotes.

Ornate Flycatcher *Myiobrycon ornatus*

White-eyed Tody-Tyrant *Hemitriccus zosterops*

Pearly-vented Tody-Tyrant *Hemitriccus margaritaceiventer* Non leader.

Black-throated Tody-Tyrant *Hemitriccus granadensis* Heard.

Southern Bentbill ♦ *Oncostoma olivaceum* Heard.

Scale-crested Pygmy Tyrant *Lophotriccus pileatus*

Pale-eyed Pygmy Tyrant *Atalotriccus pilaris*

Rufous-crowned Tody-Flycatcher* *Poecilotriccus ruficeps*

Slaty-headed Tody-Flycatcher *Poecilotriccus sylvia*

Common Tody-Flycatcher *Todirostrum cinereum*

Yellow-browed Tody-Flycatcher *Todirostrum chrysocrotaphum*

Black-headed Tody-Flycatcher *Todirostrum nigriceps* Seen below the Cerulean Reserve.

Western Olivaceous Flatbill *Rhynchocyclus aequinoctialis*

Yellow-olive Flatbill (Y-o Flycatcher) *Tolmomyias sulphurescens*

Grey-crowned Flatbill (E) *Tolmomyias poliocephalus*

Ochre-lore Flatbill (O-I Flycatcher) *Tolmomyias flaviventris*

White-throated Spadebill *Platyrinchus mystaceus* Heard.

Cinnamon Flycatcher *Pyrrhomyias cinnamomeus*

Black-billed Flycatcher ♦ *Aphanotriccus audax* Seen well at Paujil

Fuscous Flycatcher ♦ *Cnemotriccus fuscatus* The cabanisi race was seen at Los Besotes and the duidae (Campina Flycatcher) race was seen at Mitu.

Black Phoebe *Sayornis nigricans*

Olive-sided Flycatcher *Contopus cooperi* A couple were seen at Mitu

Smoke-colored Pewee *Contopus fumigatus*

Eastern Wood Pewee *Contopus virens*

Acadian Flycatcher *Empidonax virescens*

Vermilion Flycatcher *Pyrocephalus obscurus*

Streak-throated Bush Tyrant *Myiotheretes striaticollis*

White-headed Marsh Tyrant *Arundinicola leucocephala*

Yellow-bellied Chat-Tyrant *Silvicultrix diadema*

Slaty-backed Chat-Tyrant *Ochthoeca cinnamomeiventris* Heard.

Rufous-breasted Chat-Tyrant *Ochthoeca rufipectoralis*

Brown-backed Chat-Tyrant *Ochthoeca fumicolor*

Cattle Tyrant *Machetornis rixosa*

Piratic Flycatcher *Legatus leucophaeus*

Rusty-margined Flycatcher *Myiozetetes cayanensis*

Dusky-chested Flycatcher *Myiozetetes luteiventris*

Great Kiskadee *Pitangus sulphuratus*

Yellow-throated Flycatcher *Conopias parvus* Seen well at Mitu.

Golden-bellied Flycatcher *Myiodynastes hemichrysus*

Streaked Flycatcher *Myiodynastes maculatus*

Boat-billed Flycatcher *Megarynchus pitangua*

Sulphury Flycatcher *Tyrannopsis sulphurea*

Tropical Kingbird *Tyrannus melancholicus*

Fork-tailed Flycatcher *Tyrannus savanna*

Rufous Mourner *Rhytiperna holerythra* Seen at Paujil and the Piha Reserve.

Dusky-capped Flycatcher *Myiarchus tuberculifer*

Venezuelan Flycatcher ◊ *Myiarchus venezuelensis* Seen at Los Besotes and above La Jagua.

Short-crested Flycatcher *Myiarchus ferox*

Apical Flycatcher ◊ *Myiarchus apicalis* Endemic. Seen well in the Chicamocha canyon.

Pale-edged Flycatcher *Myiarchus cephalotes*

Great Crested Flycatcher *Myiarchus crinitus*

Brown-crested Flycatcher *Myiarchus tyrannulus*

Rufous-tailed Flatbill *Ramphotrigon ruficauda*

Citron-bellied Attila ◊ *Attila citriniventris* Seen well at Mitu.

Bright-rumped Attila (Flammulated A) *Attila spadiceus*

Bright-rumped Attila *Attila [spadiceus] spadiceus* Heard.

Scaled Fruiteater *Ampelioides tschudii* Heard.

Guianan Cock-of-the-rock ◊ *Rupicola rupicola* Heard.

Red-crested Cotinga *Ampelion rubrocristatus*

Screaming Piha *Lipaugus vociferans* Heard.

Chestnut-capped Piha ◊ *Lipaugus weberi* Endemic. Seen well at the Piha Reserve.

Spangled Cotinga *Cotinga cayana*

Purple-breasted Cotinga *Cotinga cotinga* A male was seen briefly at Mitu.

Pompadour Cotinga ♦ *Xipholena punicea* A female was seen at Mitu.

Dwarf Tyrant-Manakin *Tyranneutes stolzmanni* Heard.

Saffron-crested Tyrant-Manakin ♦ *Neopelma chrysocephalum* Seen well at Mitu.

Yellow-headed Manakin ♦ *Chloropipo flavicapilla* Endemic. Seen well at La Romera.

Lance-tailed Manakin *Chiroxiphia lanceolata* Heard.

Golden-winged Manakin ♦ *Masius chrysopterus*

White-bibbed Manakin ♦ *Corapipo leucorrhoea* A brief male below the Piha Reserve.

Black Manakin ♦ *Xenopipo atronitens* Seen well at Mitu.

Blue-capped Manakin *Lepidothrix coronata*

Yellow-crested Manakin ♦ *Heterocercus flavivertex* Seen well at Mitu.

White-bearded Manakin *Manacus manacus*

Golden-collared Manakin *Manacus vitellinus* Seen in the Cauca valley.

Striolated Manakin ♦ *Machaeropterus striolatus*

White-crowned Manakin *Pseudopipra pipra* Seen at the Piha Reserve and Mitu. More than one cryptic species may be involved.

Golden-headed Manakin *Ceratopipra erythrocephala*

Sulphur-rumped Myiobius *Myiobius sulphureipygius*

Whiskered Myiobius *Myiobius barbatus*

Black-crowned Tityra *Tityra inquisitor*

Black-tailed Tityra *Tityra cayana*

Russet-winged Schiffornis ♦ *Schiffornis stenorhyncha* Heard.

Brown-winged Schiffornis *Schiffornis turdina* Seen at Mitu.

White-browed Purpletuft *Iodopleura isabellae* Seen well at Mitu.

Cinereous Becard *Pachyramphus rufus*

White-winged Becard *Pachyramphus polychopterus*

Black-capped Becard *Pachyramphus marginatus* Heard.

Pink-throated Becard *Pachyramphus minor*

Rufous-browed Peppershrike *Cyclarhis gujanensis*

Black-billed Peppershrike ♦ *Cyclarhis nigrirostris*

Yellow-browed Shrike-Vireo ♦ *Vireolanius eximius* Fantastic views below San Vicente.

Brown-headed Greenlet ♦ *Hylophilus brunneiceps* Seen well at Mitu.

Lemon-chested Greenlet *Hylophilus thoracicus* Heard.

Scrub Greenlet *Hylophilus flavipes*

Rufous-naped Greenlet ◊ *Pachysylvia semibrunnea*

Chivi Vireo *Vireo chivi*

Brown-capped Vireo *Vireo leucophrys*

Yellow-throated Vireo ◊ *Vireo flavifrons*

Black-collared Jay *Cyanolyca armillata* Seen in both the Western and Eastern Andes.

Black-chested Jay *Cyanocorax affinis*

Azure-naped Jay ◊ *Cyanocorax heilprini* Seen well at Mitu.

Inca Jay *Cyanocorax yncas*

White-winged Swallow *Tachycineta albiventer*

White-banded Swallow *Atticora fasciata*

Blue-and-white Swallow *Pygochelidon cyanoleuca*

Black-collared Swallow *Pygochelidon melanoleuca* Seen well at Mitu.

Southern Rough-winged Swallow *Stelgidopteryx ruficollis*

Grey-breasted Martin *Progne chalybea*

Barn Swallow *Hirundo rustica*

Black-capped Donacobius *Donacobius atricapilla* Heard.

Bicolored Wren *Campylorhynchus griseus*

Grass Wren *Cistothorus platensis* Heard.

Sooty-headed Wren ◊ *Pheugopedius spadix* Seen well at the Piha Reserve.

Black-bellied Wren ◊ *Pheugopedius fasciatoventris* Seen well at Paujil.

Whiskered Wren *Pheugopedius mystacalis*

Coraya Wren *Pheugopedius coraya* Heard.

Rufous-breasted Wren *Pheugopedius rutilus*

Speckle-breasted Wren ◊ (Colombian W) *Pheugopedius sclateri* Heard.

Rufous-and-white Wren *Thryophilus rufalbus*

Antioquia Wren ◊ *Thryophilus sernai* Endemic. Seen well in the Cauca Valley.

Niceforo's Wren ◊ *Thryophilus nicefori* Endemic. Seen well in the Chicamocha Canyon.

Bay Wren *Cantorchilus nigricapillus*

House Wren (Southern H W) *Troglodytes aedon*

Mountain Wren *Troglodytes solstitialis*

White-breasted Wood Wren *Henicorhina leucosticte* Heard.

Grey-breasted Wood Wren *Henicorhina leucophrys*

Southern Nightingale-Wren (Scaly-breasted W) *Microcerculus marginatus* Heard.

Trilling Gnatwren *Ramphocaenus melanurus*

Collared Gnatwren ◊ *Microbates collaris* Seen well at Mitu.

Tropical Gnatcatcher *Polioptila plumbea*

Tropical Mockingbird *Mimus gilvus*

Andean Solitaire *Myadestes ralloides*

Orange-billed Nightingale-Thrush *Catharus aurantiirostris*

Swainson's Thrush *Catharus ustulatus*

Chestnut-bellied Thrush ◊ *Turdus fulviventris* Seen well at the Bushbird Reserve.

Glossy-black Thrush *Turdus serranus*

Great Thrush *Turdus fuscater*

Lawrence's Thrush *Turdus lawrencii* Heard.

Black-billed Thrush *Turdus ignobilis*

White-necked Thrush *Turdus albicollis* Heard.

Pale-vented Thrush *Turdus obsoletus* This rare species was seen well at the Piha Reserve.

Pale-breasted Thrush *Turdus leucomelas*

Clay-coloured Thrush *Turdus grayi*

Lesser Goldfinch *Spinus psaltria*

Yellow-bellied Siskin* *Spinus xanthogastrus*

Golden-rumped Euphonia *Chlorophonia cyanocephala* Seen below the Cerluean Warbler Reserve.

Chestnut-breasted Chlorophonia ◊ *Chlorophonia pyrrhophrys* Seen in the Perija.

Plumbeous Euphonia ◊ *Euphonia plumbea* Seen well at Mitu.

White-lored Euphonia *Euphonia chrysopasta*

White-vented Euphonia *Euphonia minuta*

Thick-billed Euphonia *Euphonia laniirostris*

Fulvous-vented Euphonia ◊ *Euphonia fulvicrissa*

Orange-bellied Euphonia *Euphonia xanthogaster*

Rufous-bellied Euphonia *Euphonia rufiventris*

Yellow-throated Chlorospingus *Chlorospinus flavigularis*

Ashy-throated Chlorospingus *Chlorospingus canigularis*

Common Chlorospingus *Chlorospingus flavopectus* Seen at several sites, the ponis subspecies that was seen in the Perija is a potential split.

Yellow-browed Sparrow *Ammodramus aurifrons*

Black-striped Sparrow *Arremonops conirostris* A pair seen in the Cauca valley.

Perija Brushfinch ♦ *Arremon perijanus* A skulking individual seen in the Perija foothills.

Black-headed Brushfinch ♦ *Arremon atricapillus* Seen well near the Cerulean Warbler Reserve.

Grey-browed Brushfinch *Arremon assimilis*

Golden-winged Sparrow ♦ *Arremon schlegeli* Seen well in the Perija foothills.

Chestnut-capped Brushfinch *Arremon brunneinucha*

Rufous-collared Sparrow *Zonotrichia capensis*

White-naped Brushfinch (Yellow-throated B F) *Atlapetes [albinucha] gutturalis*

Moustached Brushfinch ♦ *Atlapetes albofrenatus* Seen well at the Bushbird Reserve.

Ochre-breasted Brushfinch ♦ *Atlapetes semirufus* Seen well at Finca La Herrera.

Slaty Brushfinch *Atlapetes schistaceus* Seen on various days. The fumidis race in the Perija is rather distinctive.

Antioquia Brushfinch ♦ *Atlapetes blancae* Endemic. Seen well at the 'new' Antpitta Reserve.

Yellow-breasted Brushfinch (Northern Rufous-naped B-F) *Atlapetes latinuchus* Non leader.

Black-fronted Brushfinch ♦ *Atlapetes nigrifrons* Seen commonly in the Perija.

Eastern Meadowlark *Sturnella magna*

Red-breasted Blackbird *Leistes militaris*

Russet-backed Oropendola *Psarocolius angustifrons*

Crested Oropendola *Psarocolius decumanus*

Olive Oropendola *Psarocolius bifasciatus*

Yellow-rumped Cacique *Cacicus cela*

Subtropical Cacique *Cacicus uropygialis* Seen well at the Cerulean Reserve.

Mountain Cacique ♦ (Northern M C) *Cacicus [chrysonotus] leucoramphus* Seen well at Soata.

Yellow-backed Oriole *Icterus chrysater*

Yellow Oriole *Icterus nigrogularis*

Baltimore Oriole *Icterus galbula*

Orange-crowned Oriole *Icterus auricapillus*

Epaulet Oriole *Icterus cayanensis*

Giant Cowbird *Molothrus oryzivorus*

Shiny Cowbird *Molothrus bonariensis*

Carib Grackle *Quiscalus lugubris*

Red-bellied Grackle ◊ *Hypopyrrhus pyrohypogaster* Endemic. Seen well at La Romera.

Colombian Mountain Grackle ◊ *Macroagelaius subalaris* Endemic Eventually we found a number of flocks at Soata, with the total thought to be forty eight individuals!

Northern Waterthrush *Parkesia noveboracensis*

Golden-winged Warbler *Vermivora chrysoptera* A single in the forest at the Cerulean Reserve.

Black-and-white Warbler *Mniotilta varia*

Tennessee Warbler *Leiothlypis peregrina*

Mourning Warbler *Geothlypis philadelphia*

American Redstart *Setophaga ruticilla*

Cerulean Warbler *Setophaga cerulean* Six seen in one loose flock at the Cerulean Reserve!

Tropical Parula *Setophaga pitiauyumi*

Bay-breasted Warbler *Setophaga castanea*

Blackburnian Warbler *Setophaga fusca*

American Yellow Warbler *Setophaga aestiva*

Blackpoll Warbler *Setophaga striata*

Black-crested Warbler *Myiothlypis nigrocristata* Seen on several days. The birds in the Perija seem vocally distinct.

Buff-rumped Warbler *Myiothlypis fulvicauda*

Grey-throated Warbler ◊ *Myiothlypis cinereicollis* Seen well at the Bushbird Reserve.

Russet-crowned Warbler *Myiothlypis coronata*

Chestnut-capped Warbler *Basileuterus delatarii*

Golden-crowned Warbler *Basileuterus culicivorus*

Three-striped Warbler *Basileuterus tristriatus*

Canada Warbler *Cardellina canadensis*

Slate-throated Whitestart *Myioborus miniatus*

Golden-fronted Whitestart ◊ *Myioborus ornatus* We saw the white-spectacled ornatus form in the eastern Andes and the golden-spectacled chrysops form in the western Andes. The chrysops form has been mooted as a potential split and would be a Colombian endemic.

Dusky-faced Tanager *Mitrospingus cassinii*

Tooth-billed Tanager (Highland Hepatic T) *Piranga lutea*

Summer Tanager *Piranga rubra*

Red-hooded Tanager ◊ *Piranga rubriceps* A male was seen at Soata.

Sooty Ant Tanager ◊ *Habia gutturalis* Endemic. Seen at Paujil and heard below the Piha Reserve.

Rose-breasted Grosbeak *Pheucticus ludovicianus*

Blue-black Grosbeak *Cyanoloxia cyanooides*

Amazonian Grosbeak ◊ *Cyanoloxia rothschildii*

Plushcap (Plush-capped Finch) *Catamblyrhynchus diadema* A couple were seen in the Perija.

Green Honeycreeper *Chlorophanes spiza*

Scarlet-and-white Tanager ◊ *Chrysothlypis salmomi* A group of three showed well at the Piha Reserve.

Yellow-backed Tanager *Hemithraupis flavicollis*

Guira Tanager *Hemithraupis guira*

Swallow Tanager *Tersina viridis*

Purple Honeycreeper *Cyanerpes caeruleus*

Red-legged Honeycreeper *Cyanerpes cyaneus*

Blue Dacnis *Dacnis cayana*

Yellow-bellied Dacnis *Dacnis flaviventer*

Turquoise Dacnis ◊ (T D-Tanager) *Dacnis hartlaubi* Endemic. Seen well at the Cerulean Reserve.

Black-faced Dacnis *Dacnis lineata*

Yellow-tufted Dacnis ◊ *Dacnis egregia*

Olive-grey Saltator *Saltator olivascens*

Streaked Saltator *Saltator striatipectus*

Buff-throated Saltator *Saltator maximus*

Black-winged Saltator *Saltator atripennis*

Slate-colored Grosbeak *Saltator grossus*

Bananaquit *Coereba flaveola*

Yellow-faced Grassquit *Tiaris olivaceus*

Blue-black Grassquit *Volatinia jacarina*

White-shouldered Tanager *Loriotus luctuosus*

Fulvous-crested Tanager *Tachyphonus surinamus*

Tawny-crested Tanager *Tachyphonus delatrii*

Flame-rumped Tanager ◊ *Ramphocelus flammigerus*

Lemon-rumped Tanager ◊ (Yellow-r T) *Ramphocelus icteronotus*

Crimson-backed Tanager ◊ *Ramphocelus dimidiatus*

Silver-beaked Tanager *Ramphocelus carbo*

Grey Seedeater *Sporophila intermedia*

Yellow-bellied Seedeater *Sporophila nigricollis*

Thick-billed Seed Finch *Sporophila funerea*

Chestnut-bellied Seed Finch *Sporophila angolensis*

Chestnut-bellied Seedeater *Sporophila castaneiventris*

Ruddy-breasted Seedeater *Sporophila minuta*

Oleaginous Hemispingus *Sphenopsis frontalis* Seen in the Perija and La Romera.

White-eared Conebill ♦ *Conirostrum leucogenys* Seen below the Cerulean Reserve.

Blue-backed Conebill *Conirostrum sitticolor*

Saffron Finch *Sicalis flaveola*

Bluish Flowerpiercer *Diglossa caerulescens*

Masked Flowerpiercer *Diglossa cyanea*

Indigo Flowerpiercer ♦ *Diglossa indigotica* Seen at the Piha Reserve.

Rusty Flowerpiercer *Diglossa sittoides*

Chestnut-bellied Flowerpiercer ♦ *Diglossa gloriosissima* Endemic. Seen well at Colibri del Sol.

White-sided Flowerpiercer *Diglossa albilatera*

Black-throated Flowerpiercer *Diglossa brunneiventris* The isolated vuilleumieri race was seen at the 'new' Antpitta Reserve. A potential split.

Black Flowerpiercer *Diglossa humeralis*

Purplish-mantled Tanager ♦ *Iridosornis porphyrocephalus* Seen at the Piha Reserve.

Fawn-breasted Tanager *Pipraeidea melanonota*

Buff-breasted Mountain Tanager *Dubusia taeniata*

Hooded Mountain Tanager *Buthraupis montana*

Blue-capped Tanager *Sporathraupis cyanocephala*

Blue-winged Mountain Tanager *Anisognathus somptuosus*

Scarlet-bellied Mountain Tanager *Anisognathus igniventris*

Lacrimose Mountain Tanager *Anisognathus lacrymosus* Seen in the Perija and the Andes. The pale pallidorsalis race in the Perija is a potential split.

Black-and-gold Tanager ♦ *Bangsia melanochlamys* Endemic. Seen well at the Piha Reserve.

Magpie Tanager *Cissopis leverianus*

Black-faced Tanager *Schistochlamys melanopis*

Speckled Tanager *Ixothraupis guttata*

Blue-grey Tanager *Thraupis episcopus*

Glaucous Tanager ♦ *Thraupis glaucocolpa* Seen at Los Besotes.

Palm Tanager *Thraupis palmarum*

Black-headed Tanager* *Stilpnia cyanoptera*

Black-capped Tanager *Stilpnia heinei*

Golden-hooded Tanager *Stilpnia larvata*

Blue-necked Tanager *Stilpnia cyanicollis*

Masked Tanager *Stilpnia nigrocincta*

Scrub Tanager ◊ *Stilpnia vitriolina*

Blue-and-black Tanager *Tangara vassorii*

Beryl-spangled Tanager *Tangara nigroviridis*

Metallic-green Tanager *Tangara labradorides*

Bay-headed Tanager *Tangara gyrola*

Saffron-crowned Tanager *Tangara xanthocephala* Non leader.

Golden Tanager *Tangara arthus*

Silver-throated Tanager *Tangara icterocephala*

Plain-colored Tanager *Tangara inornata*

Turquoise Tanager *Tangara mexicana*

Paradise Tanager *Tangara chilensis*

Opal-rumped Tanager *Tangara velia*

MAMMALS

Common Opposum *Didelphis marsupialis* Seen at Mitu.

Crab-eating Fox *Cerdocyon thous*

Humboldt's White-fronted Capuchin *Cebus albifrons* Seen at Paujil.

Yellow-handed Titi *Cheracebus Lucifer* Seen at Mitu.

Mottled-face Tamarin *Saguinus inustus* Heard at Mitu.

White-footed Tamarin *Saguinus leucopus* Seen below the Piha Reserve.

Colombian Red Howler *Alouatta seniculus* Heard at several sites.

Variegated Spider Monkey ◊ *Ateles hybridus* Seen well at Paujil.

Central American Agouti *Dasyprocta punctata*

Neotropical Pygmy Squirrel *Sciurillus pusillus* Seen at Mitu.

Western Dwarf Squirrel *Microsciurus mimulus*

Red-tailed Squirrel (Tropical Red S) *Sciurus granatensis*



White-tipped Dove (image by Trevor Ellery)



Swainson's Thrush (image by Trevor Ellery)



Saffron-crested Tyrant-Manakin (image by Trevor Ellery)



Perija Parakeet (image by Trevor Ellery)



Ochre-breasted Antpitta (image by Trevor Ellery)



Inca Jay (image by Trevor Ellery)



Coopmans's Tyrannulet (image by Trevor Ellery)



Chestnut-bellied Thrush (image by Trevor Ellery)



Black-and-gold Tanager (image by Trevor Ellery)



Apical Flycatcher (image by Trevor Ellery)



Antioquia Wren (image by Trevor Ellery)



Black Manakin (image by Trevor Ellery)



Chestnut-bellied Flowerpiercer (image by Trevor Ellery)



Colombian Mountain Grackle (image by Trevor Ellery)



Grey-throated Warbler (image by Trevor Ellery)



Mountain Wren (image by Trevor Ellery)



Perija Metal-tail (image by Trevor Ellery)



Rufous-breasted Chat-Tyrant (image by Trevor Ellery)



Streak-headed Woodcreeper (image by Trevor Ellery)



White-fronted Nunbird (image by Trevor Ellery)



White-chested Puffbird (image by Trevor Ellery)



White-crowned Manakin (image by Trevor Ellery)



Spectacled Owl (image by Trevor Ellery)



Sooty Ant Tanager (image by Trevor Ellery)



Recurve-billed Bushbird (image by Trevor Ellery)



Paisa Antpitta (image by Trevor Ellery)



Maroon-tailed Parakeet (image by Trevor Ellery)



Choco Screech Owl (image by Trevor Ellery)



Forest Elaenia (image by Trevor Ellery)



Blue-billed Curassow (image by Trevor Ellery)



Black Bushbird (image by Trevor Ellery)



Yellow-browed Shrike-Vireo (image by Trevor Ellery)



Tropical Parula (image by Trevor Ellery)



Sword-billed Hummingbird (image by Trevor Ellery)



Slaty Antwren (image by Trevor Ellery)



Sickle-winged Guan (image by Trevor Ellery)



Red-bellied Grackle (image by Trevor Ellery)



Perija Thisletail (image by Trevor Ellery)



Perija Tapaculo (image by Trevor Ellery)



Paramillo Tapaculo (image by Trevor Ellery)




Orange-cheeked Parrot (image by Trevor Ellery)



Ochre-breasted Brushfinch (image by Trevor Ellery)



Mottled Owl (image by Trevor Ellery)



Magdalena Antbird (image by Trevor Ellery)



Laughing Falcon (image by Trevor Ellery)



Grey-bellied Antbird (image by Trevor Ellery)



Dusky-billed Parrotlet (image by Trevor Ellery)



Cundinamarca Antpitta (image by Trevor Ellery)



Cinnamon Screech Owl (image by Trevor Ellery)



Chestnut-capped Piha (image by Trevor Ellery)



Chestnut-capped Brushfinch (image by Trevor Ellery)



Chestnut-backed Antbird (image by Trevor Ellery)



Black-headed Antbird (image by Trevor Ellery)



Black-fronted Brushfinch (image by Trevor Ellery)



Black Caracara (image by Trevor Ellery)



Barred Puffbird (image by Trevor Ellery)



Azure-naped Jay (image by Trevor Ellery)